

PCN Insights Pack 2024 Stevenage South

Stefania Mistretta, PHM Champion Hannes van der Merwe, PHM Champion Jaron Inward, Snr PHM Champion

Working together for a healthier future



Introduction

This latest version of the PCN pack has been developed around the ICB Clinical Priorities signed off by the ICB Board in March 23 and how they align to the population outcomes we are aiming to achieve. (See tables on the right hand side.)

The data contained within this pack compares the PCN data against Place and ICB. An overview table of all the data is available on the following page with areas of opportunity highlighted.

Some data will not be available at PCN level e.g. mortality rate for CVD. Where this is the case, proxy measures that will lead to improving this will be included e.g. early identification.

Where opportunities link with areas within the ECF or QOF a selection of related indicators have been shared alongside a link to Ardens Manager.

The data within this pack are shown at PCN level and are health focused. However to improve outcomes for our population, input from many partners is required. This pack can facilitate discussion within your Integrated Neighbourhood Teams as described within the Primary Care Strategy and also the wider HCP transformation spaces supported by your Primary Care Transformation leads.

Packs for other PCNs, Localities and Place can be found:

<u>Population health management – Hertfordshire and West Essex Integrated Care System</u> (hertsandwestessexics.org.uk)

Here you can also find previous PCN packs outlining the descriptive demographics for the PCNs.

Area	Clinical Priority
СҮР	Improved Readiness for school in children eligible for FSM Reduce rates of Childhood obesity Reduced unnecessary A&E attendances and admissions
Prevention and Health Inequalities	Reduced premature mortality rate for CVD
LTC & Frailty	Reduce attendance and admissions for falls, people with frailty and people in last year of life Development of more proactive, preventative care models for management of LTC and Frailty
Mental Health	Reducing suicide rates and attendances/ admission rates for self-harm Reducing rates of A&E attendances involving substance misuse and violence

Outcome	Definition
Improve life expectancy	Average age at death for people who have died in the last 12 months
Improve healthy life expectancy	Average age of people who have left the 'healthy' segment in the last 12 months
Reduce the proportion of people living with advanced disease and complexity	Proportion of the registered population who are in the advanced disease and complexity segment (segment)
Reduce the rate of ambulatory care sensitive emergency hospital admissions	Rate of ambulatory care sensitive emergency admissions in the last 12 months
Reduce the overall spend on emergency hospital admissions	Spend on Emergency hospital admissions within a financial year





Stevenage South at a Glance

The table on the right provides a summary of the data contained within this pack. It highlights how the PCN data compares with Place and ICB. For Stevenage South areas of opportunity highlighted are:

- Childhood obesity reception and Year 6
- A&E attendance for asthma in children
- Observed versus expected prevalence of LTC
- Annual Reviews completed for LTCs
- Control of hypertension
- Admissions for Chronic Ambulatory Care Sensitive Conditions
- Admissions for falls and hip fractures in the over 75s
- Admissions for self-harm

The following link takes you to Ardens Manager where there are reports. Here you will find the latest information on identification of LTCs and details of case finding Ardens searches available within EMIS and Systmone.

https://app.ardensmanager.com/login


e	Clinical Priority	Metric Metric	PCN compared to Place average	PCN compared to ICB average
	Childhood obesity	% of children in Reception who are overweight	↑	↑
	Cilianood obesity	% of children in Year 6 who are overweight	↑	↑
		A&E Attendances for Asthma (Children)	↑	↑
	Reduce rates of	Admissions for Asthma (Children)	\	\
	emergency care for children and young	Admissions for Wheeze (Children)	↑	\leftrightarrow
	people	Admissions for Diabetes (Children)	\	\
		Admissions for Epilepsy (Children)	\leftrightarrow	↑
		Lifestyle risk factors: Smoking	↑	↑
		Observed versus expected prevalence	\	\
	Prevention and health	Annual Reviews completed for LTCs	\	\
	inequalities (Premature mortality for CVD)	% of people with AF treated with Anti Coagulant	\leftrightarrow	\leftrightarrow
	, , ,	Control of hypertension	V	\
	Preventative, Proactive	Identification of hypertension	\leftrightarrow	\leftrightarrow
	care models for LTC	% of people for secondary prevention CVD who are on low and medium $$ intensity statins	\leftrightarrow	\leftrightarrow
		% of diabetics with all 8 care processes completed	↑	\leftrightarrow
		Admissions for ACS conditions	↑	↑
	Preventative, Proactive	Admissions for falls (75+)	↑	↑
:	care models for frailty and EOL	Admissions for Hip Fractures (75+)	↑	↑
	Mantal Harlth	Prevalence of Mental Health Conditions including LD	\leftrightarrow	\leftrightarrow
	Mental Health	Admissions for Self-Harm	↑	↑


Rates of Childhood Obesity

CYP outcome - Every child will have the best start and live a healthy life

ICB overarching outcome of **Improving Healthy life** expectancy

- In keeping with the national trend, the PCN rates for Childhood Obesity are higher for year 6 in comparison to reception children.
- Compared to the ICB and Place rate, Stevenage South PCN has a slightly higher rate of Childhood Obesity for Children in Reception. The rate of Childhood Obesity at Year 6 is higher when comparing with Place and the ICB.
- The data suggest that there is a deterioration from reception to Year 6 in childhood obesity in the PCN position against Place and ICB.











A&E attendances for Asthma (CYP)

CYP outcome – Reduce the number of unplanned admissions for long term conditions

- Compared to Place and the overall ICB, Stevenage South PCN has a slightly higher rate of A&E attendances for Children and Young People with Asthma (rolling years data on the right-hand side).
- Rates of Children and Young People attending A&E for Asthma have remained fairly steady since 20/21.









Admissions for Asthma (CYP)


- Compared to Place and the ICB, Stevenage South PCN has a lower rate of children's admissions for Asthma (rolling years data on the right-hand side). It is to be noted there is no data for two of the PCN's practices.
- The trend data shows a decreasing trend for the PCN, however the numbers are small which can cause fluctuation in rates.
- Lower Proportion of Asthma Reviews are carried out within Stevenage South PCN in comparison to Place and the ICB. However, the QOF is for all ages and children specific reviews cannot be identified within the data.



CYP outcome – Reduce the number of unplanned admissions for long term conditions

ICB overarching outcome of reduce the proportion of people living with advanced disease and complexity







Source: SUS; QOF

Admissions for Wheeze (CYP)

CYP outcome – Reduce the number of unplanned admissions for long term conditions

- Stevenage South PCN has a higher rates of Children and Young People admitted to hospital for abnormalities of breathing including wheeze compared to Place and similar to the ICB. Recent analysis of UEC data showed Wheeze as a significant reason for admission within young children across HWE.
- Rates of Children and Young People admitted to Hospital for Wheeze fluctuate annually with the latest forecast outturn from November data showing a slight decrease on the previous year.
- When looking at the data by practice there is variation between the practices.









Admissions for Diabetes (CYP)

CYP outcome – Reduce the number of unplanned admissions for long term conditions

- Diabetes is identified as an area of focus within the Core 20 plus 5 for children. The PCN rate for admissions is lower than both place and ICB.
- The numbers of children admitted for diabetes are small and this should be considered when looking at the data. There were no admissions for the Knebworth & Marymead Practice within the 12 months up to November 23.
- The data for diabetes will continue to be monitored at wider HCP and ICB footprints.









Admissions for Epilepsy (CYP)

CYP outcome – Reduce the number of unplanned admissions for long term conditions

- Epilepsy is identified as an area of focus within the Core 20 plus 5 for children. The PCN rate for admissions is higher than ICB and similar to place.
- The numbers of children admitted for epilepsy are small and therefore fluctuations in the trend are more prominent. When looking at the data by practice, the Bedwell Medical Centre have a higher rate of Children and Young People admitted to hospital compared to the other practices.
- The data for epilepsy will continue to be monitored at wider HCP and ICB footprints.










Prevention and health inequalities – Lifestyle factors - Smoking

- Stevenage South PCN data for smoking shows a similar picture to the Place and ICB for smoking status recorded.
- A higher proportion of patients have been offered treatment for smoking compared to ICB and place.
- The table to the right gives detail by condition of the opportunity for further recording of smoking status.
 This shows the position in January. The latest position can be found on https://app.ardensmanager.com/login

	ECF 2023-24 - Condition Section Under Smoker, Smoking Status, and Smoking Status Recorded - as of Jan. 2024											
	Pre-Di	abetes	Diab	etes	Atrial Fibrillation							
	Remaining % of	Smoking Available	Remaining % of	Smoking Available	Remaining % of	Smoking Available						
Practices	Population with a	Patients - Number	Population with a	Patients - Number	Population with a	Patients - Number						
	Smoking status		Smoking status		Smoking status							
Bedwell Medical Centre	31%	765	11%	1044	0%	22						
King George Surgery	43%	1270	11%	1418	0%	20						
Knebworth & Marymead Me	47%	306	41%	853	0%	14						
Shephall Way Surgery	36%	522	16%	663	0%	12						







Source: Link: QOF Data Set & ECF Jan. 2024





Prevention and health inequalities Early Identification: Expected vs observed prevalence

The data on here shows the national modelled estimated prevalence for the PCN compared with the latest published QOF prevalence for the PCN.


- Stevenage South PCN recorded prevalence compared with the modelled estimated prevalence for the PCN is lower across all conditions.
- The data shows an opportunity for further identification for these conditions. Case finding searches can be found within the Ardens Suite of searches: https://app.ardensmanager.com/login

Disease Detection Modelling for Stevenage
South PCN - No. of New Diagnoses to Meet ICS
& PLACE Rates - 2023/24

Disease / Canadisian	Number to meet	Number to meet
Disease/ Condition	ICS rate	PLACE rate
Asthma	3377	43
Atrial Fibrillation		129
Chronic Kidney Disease	1569	
Chronic Obstructive Pulmonary Disease		58
Coronary Heart Disease	1573	72
Diabetes Mellitus		
Epilepsy	327	
Heart Failure		23
Hypertension	8011	254
Non-Diabetic Hyperglycaemia		331
Peripheral Arterial Disease	252	0
Stroke and Transient Ischaemic Attack	954	42

LTC Outcome – Proportion of people with a long-term condition who feel able to manage their condition

ICB overarching outcome of reduce the proportion of people living with advanced disease and complexity







Development of more proactive, preventative care models for LTC - Prevalence

LTC Outcome – Reduce the proportion of people with a long-term condition who are in the advanced disease & complexity or frailty & end of life segments

Stevenage South BCN - Long-Term Conditions 2022-23 OOE Prevalence with 3 Year Trend

ICB overarching outcome of reduce the proportion of people living with advanced disease and complexity

- Following the expected and observed prevalence modelling in the previous slide, where there are opportunities for further identification, this slide looks at the recorded prevalence by practice for the PCN compared with Place and ICB.
- For LTCs, QOF and ECF data can help us understand areas of opportunity for improving outcomes for our population living with LTCs.

					Stevenage South PCN - Long-Term Conditions 2022-23 QOF Prevalence, with 3 Year Trend.						
	QOF 22-	QOF 22-	QOF 22-	KNEBW	KNEBWORTH &		SHEPHALL HEALTH		KING GEORGE &		MEDICAL
OOF Disease / Condition	23 -	23 -	23 -	MARYMEA	PRACTICE	CEN	TRE	MANOR	HOUSE	CENTRE	
QOF Disease/ Condition	ICB %	PLACE %	PCN %	QOF 2022-	3 Year	QOF 2022-	3 Year	QOF 2022-	3 Year	QOF 2022-	3 Year
				23	Trend	23	Trend	23	Trend	23	Trend
Asthma	6.16%	6.40%	6.43%	6.49%		6.56%		6.61%		6.06%	
Atrial fibrillation	2.09%	2.12%	2.00%	2.51%	_	1.83%		1.77%	\	1.90%	_/
Chronic kidney disease	3.46%	2.94%	3.25%	3.18%	_	2.93%		2.99%	_	3.88%	
Chronic obstructive pulmonary disease (COPD)	1.49%	1.55%	1.78%	1.91%		2.11%		1.27%	_	1.83%	
Diabetes mellitus	6.63%	6.54%	7.32%	6.79%	_/	7.21%		6.70%	_/	8.57%	/
Epilepsy	0.70%	0.73%	0.92%	0.97%	/	1.01%		0.76%	_	0.94%	/
Heart Failure	0.80%	0.74%	0.71%	0.76%	_/	0.61%	_/	0.71%	\	0.77%	_/
Hypertension	13.84%	13.83%	13.83%	14.29%	_	13.83%		12.49%	_	14.71%	_/
Non-diabetic hyperglycaemia	6.42%	5.29%	5.67%	2.34%		6.48%	/	6.71%	\	7.16%	
Peripheral arterial disease	0.44%	0.45%	0.46%	0.47%		0.45%		0.44%	_	0.49%	\
Secondary prevention of coronary heart disease	2.67%	2.63%	2.59%	2.93%		2.47%		2.41%	_	2.54%	_
Stroke and transient ischaemic attack	1.63%	1.71%	1.75%	2.13%	_	1.50%	_	1.62%	\	1.76%	~/



Development of more proactive, preventative care models for LTC : Annual Reviews

- The table on the right shows a summary of the percentage of patients receiving an annual review or risk assessment by condition.
- Where the cell is highlighted the percentage is lower than the Place value.
- The data shows that the PCN practices have lower percentages for all areas apart from heart failure.
- The source of data in this table is QOF national reporting. More detailed information with the latest position is available to practices via https://app.ardensmanager.com/login


	ICB	E&N	Stevenage South PCN	Bedwell Medical Centre	King George & Manor House Surgeries	Knebworth & Marymead Practice	Shephall Health Centre
% of AF Patients with Stroke Risk Assessed in the last 12 months	92.9	91.8	87.4	93.4	84.9	81.1	94.0
The % of patients aged 45 or over who have a record of blood pressure in the preceding 5 years	85.7	85.9	84.4	86.0	84.0	84.5	82.8
The % of patients with a diagnosis of heart failure on the register, who have had a review in the preceding 12 months	72.7	67.0	69.0	95.0	63.0	55.3	65.5
The % of patients with asthma, on the register, who have had an asthma review in the preceding 12 months	64.0	65.3	60.8	85.8	54.0	51.4	59.1
The % of patients with COPD, on the register, who have had a review in the preceding 12 months	75.7	77.1	76.4	94.9	74.0	56.2	84.1
The % of patients with diabetes, on the register, with a record of a foot examination and risk classification: 1) low risk (normal sensation, palpable pulses), 2) increased risk (neuropathy or absent pulses), 3) high risk (neuropathy or absent pulses plus deformity or skin changes in previous ulcer) or 4) ulcerated foot within the preceding 12 months	75.5	76.4	76.3	70.2	87.5	68.8	71.3
The % of patients with non-diabetic hyperglycaemia who have had an HbA1c or fasting blood glucose performed in the preceding 12 months.	82.9	82.9	80.9	87.3	79.8	78.5	75.2






Prevention and health inequalities – Atrial Fibrillation

- Stevenage South PCN recorded prevalence for Atrial Fibrillation is lower than both Place and the ICB prevalence.
- Once identified the percentage of patients currently treated with anti-coagulant drug therapy is similar to the Place and ICB.
- The data suggests there is further opportunity for identification of people with AF. Case finding Ardens searches are detailed within https://app.ardensmanager.com/login









Prevention and health inequalities – Hypertension

- Stevenage South PCN recorded prevalence for hypertension is similar to both Place and the ICB prevalence.
- Once identified the data shows the percentage of patients in whom the latest BP reading is below the age-appropriate treatment threshold is lower than Place and ICB, there is variation between the practices.
- The data suggests there is further opportunity for identification of people with hypertension. Case finding Ardens searches are detailed within https://app.ardensmanager.com/login








Lipid management: Percentage of people on lipid lowering therapy for secondary prevention who are on low and medium intensity statins


- National lipid management pathways (<u>Link to guidance</u>) recommend the use of high intensity statins for all people with a history of Cardiovascular disease as well as where high dose statins are needed to control cholesterol. People on high intensity statins will see a greater reduction in c-LDL levels and reduce the risk of cardiovascular events.
- Data from OpenPrescribing provides information on the proportion of people who are on statins that are currently prescribed low or medium intensity doses. The ICS is a negative outlier in this area, with a high proportion of people not on a high intensity statin.
- The data for Stevenage South PCN shows that there is an opportunity to improve statin treatment, prescribing a higher proportion of people onto high intensity statins. The PCN is in the 57th percentile with 29.5% of people not on high intensity statins. This compares to 28.3% nationally.






Development of more proactive, preventative care models for LTC: 8 Care Processes & 3 treatment targets (all diabetes type 1 & 2)


- The percentage of people living with diabetes who have received the 8 care processes in Stevenage South PCN is similar to the ICB and higher than Place. However, for the three treatment targets the PCN data shows a lower percentage than Place and ICB.
- The latest information can be found within Ardens Manager.








Reduction in admissions rates of Chronic Ambulatory Care Sensitive (ACS) conditions



LTC Outcome – Reduce the rate of ambulatory care sensitive emergency hospital admissions

ICB overarching outcome of Reduce the rate of emergency admission for chronic ambulatory care sensitive conditions associated with LTCs



- Ambulatory care sensitive (ACS) conditions are conditions where effective community care and person-centred care can help prevent the need for hospital admission. (Nuffield Trust)
- Stevenage South PCN's admission rate for Chronic ACS conditions is higher than the ICB rate and the Place rate when looking at the 12 months data up to November 2023.
- AF & Flutter, COPD and Diseases of the blood (including diabetes), are the conditions with the highest volume and are also showing an upward trend.
- The data on page 11 looking at observed versus expected prevalence shows an opportunity for further identification of those with AF, Coronary Heart disease, COPD and Heart Failure.

	PCN Per		
Chronic Ambulatory Care Sensitive Conditions	100,000	5 Year	2024/25
for Stevenage South PCN	Rate Apr-23	Trend	Trajectory
	to Nov-23		
Angina: Angina pectoris	13.82		UP
Asthma	42.12	~	UP
Atrial fibrillation and flutter	164.09	\	UP
COPD	222.88	\	UP
Congestive heart failure	129.31	\\ \	UP
Diseases of the blood	235.99	\	UP
Epilepsy	32.21		DOWN
Hypertension	21.70	~~	UP
Mental and behavioural disorders		~	
Nutritional, endocrine and metabolic	87.52		UP


Source: SUS Link: Chronic ACS Conditions & NHSOF


Emergency Admission rates for Falls in persons aged +75

Frailty and EoL Outcome –
Decrease rates of +75s
emergency admissions for falls
within the community

ICB overarching outcome of Reduce the overall spend on emergency hospital admissions

- When looking at the rolling 12 months up to November 2023 the data shows that Stevenage South PCN has a higher rate of admissions for falls than Place and ICB.
- There is variation in the data for the practices within the PCN.
- Data in the following pages shows the data for the PCN compared with Place and PCN for areas within the ECF that aim to support reducing falls. Frailty Clinical Leads will be able to advise on current programmes of work within your area aimed at reducing falls.











Emergency admission rates for Hip fractures in all over 75's

Frailty and EoL Outcome –
Decrease rates of +75s
emergency admissions for falls
within the community

ICB overarching outcome of Reduce the overall spend on emergency hospital admissions

- The 12 months data up to November 2023 the data shows that Stevenage South PCN has a higher rate of admissions for hip fractures than Place and ICB. This follows from what is observed within the falls data.
- When looking at the data by PCN the small numbers will cause fluctuations over the years. Although the latest trend data shows a decrease for the latest year against the previous year.
- The ECF indicators on the next page shows potential areas of opportunity through benchmarking the PCN against Place and ICB.







ECF indicators for frailty and **EOL**

- The data shows that Stevenage South PCN has a slightly lower percentage of falls frat scores completed, when compared to Place but higher than ICB as at end Dec 23Where the values are zero, this may be due to the data not being submitted.
- The percentage of the population recorded as moderately or severely frail is lower than Place and ICB indicating possible opportunity for further identification.
- The data contained within the table below is up to the end of December, the latest position can be found at Ardens Manager.

Stevenage South Enhanced Commissiong Framework (ECF) 2023-24 for the period 1 April 23 to 31 Dec 23


	Frailty			EOL						
	Mod/Sev + falls Frat score done %	Mod fraily + SMR or polypharmacy medication review %	% Mod/Sev frailty of population	% population on EOL register	GSF %	DNACPR %	ACP %	PPD %	PPC %	SCR Consent %
ICB	16.0%	19.5%	1.9%	0.7%	64.9%	53.3%	11.3%	37.3%	39.8%	34.8%
ENH	20.0%	11.7%	1.7%	0.8%	77.5%	48.6%	9.3%	35.1%	37.1%	48.0%
PCN	19.0%	0.0%	0.9%	0.7%	78.6%	54.0%	4.3%	29.9%	29.4%	44.5%
Bedwell Medical Centre	0.0%	0.0%	0.0%	1.0%	93.1%	73.8%	2.3%	36.9%	34.6%	47.7%
King George & Manor House Surgeries	23.1%	0.0%	1.0%	0.4%	73.7%	55.8%	9.5%	40.0%	41.1%	67.4%
Knebworth & Marymead Practice	13.3%	0.0%	1.5%	0.7%	55.1%	53.1%	4.1%	28.6%	27.6%	39.8%
Shephall Health Centre	21.1%	0.0%	0.9%	0.8%	90.7%	18.7%	1.3%	6.7%	8.0%	16.0%




ICB overarching outcome of Improving **Healthy life expectancy**

- The data on this page looks at the recorded prevalence of conditions within the Mental Health section of QOF. It shows the PCN prevalence against the Place and ICB. Future iterations will include comparisons against modelled expected prevalence.
- The data shows that Stevenage South PCN has a similar recorded prevalence for SMI, Dementia, LD and Depression.
- The table below shows the trend over the last three years for each area.
- The following page looks at some of the wider QOF indicators around Mental Health.

	Stevenage South PCN- Mental Health Conditions 2022-2023 QOF prevalence, with 3 year trend										
	Bedwell Medic	cal Centre	Centre King George & Manor house Surgeries			h & Marymead ractice	Shephall Health Centre				
	QOF Prevalence 22-23	3 year Trend	QOF Prevalence 22-23	3 year Trend	QOF Prevalence 22-23	3 year Trend	QOF Prevalence 22-23	3 year Trend			
Dementia	0.9%		0.6%	/	1.2%	\	0.4%				
Depression	13.4%	_	14.1%		10.8%		13.7%				
Learning Disability	0.6%	_/	0.6%		0.6%		0.7%				
SMI	0.9%	_/	0.9%		0.9%		0.8%				









Mental Health QOF Indicators 22-23

- The data her shows the latest Mental Health QOF metrics for 2022-23 for SMI and Depression in comparison to Place and the ICB.
- The data shows that Stevenage South PCN have lower recorded achievement for the percentage of Patients with SMI with a care plan in comparison to Place and the ICB.
- The data contained within the table below is the latest QOF data, the latest in year position can be found at Ardens Manager.

			SMI			Depression
	% of patients with SMI who have a care plan	% of patients with SMI who have a record of BMI in the preceding 12 months	% of patients with SMI who have a record of alcohol consumption in preceding 12 months	% of patients with SMI who have a record of a lipid profile in the preceding 12 months	% of patients with SMI who have a record of blood glucose of HbA1C in preceding 12 months	% of patients with a diagnosis of depression who have been reviewed within 10-56 days
ICB	82.6	88.7	89.3	83.1	83.0	83.0
E&N	79.8	88.3	89.0	81.5	81.5	82.0
Stevenage South PCN	74.5	91.7	89.9	82.7	81.1	88.5
Bedwell Medical Centre	95.8	95.8	98.7	91.4	87.9	100.0
King George & Manor House Surgeries	92.7	97.6	97.5	97.4	97.9	85.4
Knebworth & Marymead Practice	52.1	75.0	80.2	63.2	65.9	84.8
Shephall Health Centre	57.4	100.0	81.2	76.5	68.4	84.5



Emergency Admissions Rates for Self – Harm

ICB overarching outcome of Improving Healthy life expectancy

- Stevenage South PCN has a higher rate of admissions for self harm compared with both Place and ICB.
- When looking at the data it should be noted that the numbers at PCN level are small and therefore more fluctuation between the years will be seen.
- The data will continue to be monitored at wider HCP and ICB footprints.

